

Red meat & processed meat: Summarising the Public Health isues

4 November 2015

Cristina Cleghorn, Nick Wilson, Tony Blakely



www.bigstock.com · 17926517

This blog was triggered by the recent highly

publicised review on the cancer risk from processed meat and red meat. Here we briefly look at this topic and also take a wider perspective on other aspects of meat consumption on human health and the environment, and risk communication.

The cancer agency of the World Health Organization, the International Agency for Research on Cancer (IARC), has recently <u>evaluated the carcinogenicity of consuming red and</u> <u>processed meat</u> (1). Over 400 different epidemiological studies on cancer in humans provided data on processed meat and over 700 provided data on red meat.

Processed meat

Processed meat refers to meat that has been transformed through salting, curing, fermentation, smoking, or other processes to enhance flavour or improve preservation. Most processed meats contain pork or beef, but processed meats may also contain other red meats, poultry, offal, or meat by-products such as blood. Examples of processed meat include hot dogs (frankfurters), ham, sausages, corned beef, and biltong or beef jerky as

well as canned meat and meat-based preparations and sauces.

Processed meat was classified by IARC as *carcinogenic to humans* (Group 1 – see Figure), based on *sufficient evidence* in humans that the consumption of processed meat causes colorectal cancer. Each 50 gram [g] portion of processed meat eaten daily (e.g. 1-2 rashers of bacon) was reported to increase the risk of colorectal cancer by 18% – which translates to 0.045 years of reduced non-Māori male life expectancy (or 16 days of reduced life expectancy; calculated from the BODE³ colorectal cancer screening model, personal communication <u>Dr Giorgi Kvizhinadze</u>). An association with stomach cancer was also seen, but the evidence is described as "not conclusive".

Red meat

Red meat refers to all mammalian muscle meat, including, beef, veal, pork, lamb, mutton, horse, and goat. Red meat was classified by IARC as **probably carcinogenic to humans** (Group 2A – see Figure), based on **limited evidence** that the consumption of red meat causes colorectal cancer in humans and **strong** mechanistic evidence supporting a carcinogenic effect. There is also evidence of associations with pancreatic cancer and prostate cancer.

Figure: IARC Carcinogen Classifications



What is the significance of this finding?

IARC has categorised the association of processed meat with cancer in the highest category of **strength of evidence** (see Figure), but how strongly is processed meat actually associated with cancer? Sir David Spiegelhalter, a risk professor at Cambridge University, summarised the increased risk, using the UK as an example:

"In the UK, around six out of every 100 people get bowel cancer at some point in their lives.

If they all had an extra 50g of bacon" (or hot dogs/ham/ sausages/corned beef/ beef jerky/canned meat/meat-based preparations and sauces) "a day for the rest of their lives then the risk would increase by 18% to around seven in 100 people getting bowel cancer.

"So that's one extra case of bowel cancer in all those 100 lifetime bacon-eaters" (<u>http://www.bbc.com/news/health-34615621</u>)

Only 12.7% of the NZ adult population 'Never consumed (or not consumed in past 4 weeks)' processed meat (2), and the average total consumption of processed meat (a narrower definition, ignoring corned beef, canned meat and meat-based preparations and sauces) ranged from 24g per day in Non-Māori females to 56g per day in Māori males in the last NZ adult nutrition survey (unpublished data). So these "lifetime bacon-eaters" are not too different from your average New Zealander.

A recently published meta-analysis found that those in the highest category of consumption of processed meat and red meat also had an 18% and 16% higher risk of dying from cardiovascular disease (CVD) compared to those in the lowest category of consumption (3). Similarly, the risk of type 2 diabetes was 13% higher per 100g consumption of red meat and 32% higher for 50g consumption of processed meat (4), although results from both meta-analyses should be interpreted with caution due to the potential of residual confounding. The Global Burden of Disease study estimates that approximately 841,000 total deaths per year are attributable to diets high in processed meat (5) (ranked the 14th most important risk factor for health loss in Australasia). An estimated 38,000 total deaths were attributed to red meat consumption globally each year (5).

Communicating health impacts is challenging. The public is increasingly frustrated by messages of harm from multiple food items, without clarity as to what an alternative diet is. On the other hand, research such as that summarised by IARC is essential to building the evidence base on what foods affect what health outcomes.

In the future, though, increasing emphasis needs to be placed on guidance as to a healthier diet. Of course people make trade-off decisions and some will be prepared to sacrifice (on expectation) something like several weeks of life expectancy if they prefer eating processed meat instead of healthier foods. But this life expectancy sacrifice is on average – the reality is that a small percentage of people will lose years of life expectancy, and the majority will suffer no impact. It also needs to be acknowledged, though, that we should be considering not just health impacts, but wider environmental and social impacts of a food production system.

The benefits of red meat consumption

Red meat is a good source of protein and micronutrients such as iron, zinc and B12 (6, 7).

But such nutrients can all be obtained from plant foods – and these may provide a healthier source given the lower CVD risk. For example, the evidence favouring nut consumption for reduction in CVD deaths, cancer deaths and all-cause mortality, is getting strong (see these systematic reviews (8,9)).

Should we be looking wider than human health?

Environmental impacts: Many countries around the world, including Sweden, Brazil and Italy, are starting to consider the environmental impact of food consumption when designing or updating their food and nutrition recommendations. The meat-based food system requires more energy, land, and water resources than a meatless diet (10). For example, the energy input to produce 1kcal of protein is 11 times greater for animal protein production compared to grain protein production (10). Worldwide, agricultural activity accounts for about a fifth of global greenhouse-gas emissions, 80% of this is from livestock production (11). At a time when natural resources are becoming more scarce and we need to be decreasing greenhouse gas emissions in order to meet the 350 ppm CO₂ target to avoid dangerous climate change, should countries be moving to a more sustainable sources of protein and micro-nutrients?

Ethics and animal welfare: Some people have views that it is ethical for humans to kill and eat animals, and others consider it to be unethical (there is a vast philosophical literature on the topic). But there is likely to be fairly widespread public agreement that animal welfare does matter and that farmed animals should not suffer unnecessary pain and deprivations. Unfortunately there is evidence that modern factory-farming practice causes animal suffering in various forms (see for example these two systematic reviews on pain in farmed piglets (12,13)). Reductions of processed and red meat consumption would reduce such suffering overall as long as there wasn't complete substitution onto poultry. It may also be deleterious environmentally if a reduction in meat consumption caused an increase in fish consumption (i.e. reduction of natural stocks, or aquaculture that requires feeding fish meat products as opposed to herbivore fish farming).

Conclusions

There is now strong scientific evidence of an increased risk of colorectal cancer with processed meat consumption, limited evidence of red meats being associated with colorectal cancer and some evidence of an association between red and processed meat and CVD and type 2 diabetes. Collectively all this matters since these diseases impose substantial health loss and also costs on publicly-funded health systems.

There are also other issues involved with meat production – including greenhouse gas emissions from ruminant agriculture and ethical/animal welfare concerns. Perhaps it is time for governments to more fully explore the options around reducing meat consumption at the population level – to benefit health, the financial viability of the health system, and the environment.

Authors: Dr Cristina Cleghorn, Associate Professor Nick Wilson, Professor Tony Blakely

References

1. International Agency for Research on Cancer (World Health Organisation): IARC Monographs evaluate consumption of red meat and processed meat. In., No. 240 edn. Lyon, France,; 2015: 2.

- 2. University of Otago, Ministry of Health: A Focus on Nutrition: Key findings of the 2008/09 New Zealand Adult Nutrition Survey. In. Wellington Ministry of Health; 2011.
- Abete I, Romaguera D, Vieira AR, Lopez de Munain A, Norat T: Association between total, processed, red and white meat consumption and all-cause, CVD and IHD mortality: a meta-analysis of cohort studies. British Journal of Nutrition 2014, 112(05):762-775.
- 4. Feskens EJ, Sluik D, van Woudenbergh GJ: Meat consumption, diabetes, and its complications. Current Diabetes Reports 2013, 13(2):298-306.
- Lim SS, Vos T, Flaxman AD, Danaei G, Shibuya K, Adair-Rohani H, AlMazroa MA, Amann M, Anderson HR, Andrews KG: A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010. Lancet 2013, 380(9859):2224-2260.
- 6. McAfee AJ, McSorley EM, Cuskelly GJ, Moss BW, Wallace JM, Bonham MP, Fearon AM: Red meat consumption: An overview of the risks and benefits. Meat Science 2010, 84(1):1-13.
- 7. Williams P: Nutritional composition of red meat. Nutrition & Dietetics 2007, 64(s4):S113-S119.
- Banel DK, Hu FB: Effects of walnut consumption on blood lipids and other cardiovascular risk factors: a meta-analysis and systematic review. The American Journal of Clinical Nutrition 2009, 90(1):56-63.
- 9. Grosso G, Yang J, Marventano S, Micek A, Galvano F, Kales SN: Nut consumption on all-cause, cardiovascular, and cancer mortality risk: a systematic review and metaanalysis of epidemiologic studies. The American Journal of Clinical Nutrition 2015, 101(4):783-793.
- 10. Pimentel D, Pimentel M: Sustainability of meat-based and plant-based diets and the environment. The American Journal of Clinical Nutrition 2003, 78(3):660S-663S.
- 11. McMichael AJ, Powles JW, Butler CD, Uauy R: Food, livestock production, energy, climate change, and health. Lancet 2007, 370(9594):1253-1263.
- 12. Dzikamunhenga RS, Anthony R, Coetzee J, Gould S, Johnson A, Karriker L, McKean J, Millman ST, Niekamp SR, O'Connor AM: Pain management in the neonatal piglet during routine management procedures. Part 1: a systematic review of randomized and non-randomized intervention studies. Animal Health Research Reviews / Conference of Research Workers in Animal Diseases 2014, 15(1):14-38.
- 13. O'Connor A, Anthony R, Bergamasco L, Coetzee J, Gould S, Johnson AK, Karriker LA, Marchant-Forde JN, Martineau GS, McKean J et al: Pain management in the neonatal piglet during routine management procedures. Part 2: grading the quality of evidence and the strength of recommendations. Animal Health Research Reviews / Conference of Research Workers in Animal Diseases 2014, 15(1):39-62.

Public Health Expert Briefing (ISSN 2816-1203)

Source URL:

https://www.phcc.org.nz/briefing/red-meat-processed-meat-summarising-public-health-isue s